

RESEARCH ARTICLE

Historical knowledge, richness and relative representativeness of the avifauna of the largest native urban rainforest in the world

Vinicius R. Tonetti¹, Marco A. Rego², André C. De Luca³,
 Pedro F. Develey³, Fábio Schunck⁵, Luís F. Silveira⁶

¹Departamento de Ecologia, Universidade Estadual Paulista. Avenida 24A, 1515, Jardim Bela Vista, 13506-900 Rio Claro, SP, Brazil.

²Museum of Natural Science, Louisiana State University. 70802 Baton Rouge, LA, USA.

³Avenida Pedro Paulo de Souza, 1750, Quadra HC-08. 74663-520 Goiânia, GO, Brazil.

⁴BirdLife/SAVE Brasil. Rua Fernão Dias 219, conjunto 2, 05427-010 São Paulo, SP, Brazil.

⁵Programa de Pós-Graduação, Departamento de Zoologia, Instituto de Biociências, Universidade de São Paulo. 03178-200 São Paulo, SP, Brazil.

⁶Museu de Zoologia, Universidade de São Paulo. Avenida Nazaré 481, Ipiranga, 04263-000 São Paulo, SP, Brazil.
 Corresponding author: Vinicius Rodrigues Tonetti (vrtonetti@gmail.com)

<http://zoobank.org/D811F345-6865-4000-B2F7-0E025BDADA2C>

ABSTRACT. Stretching for more than 10,000 ha in the Metropolitan Area of São Paulo, southeastern Brazil, Serra da Cantareira comprises the largest native urban rainforest in the World, harboring a rich and diverse Atlantic Forest avifauna. Despite its closeness to major urban areas, few bird surveys have been conducted there. In this article we present an updated compilation of all bird species recorded for Serra da Cantareira, including personal records from the authors. A total of 326 species have been recorded for Serra da Cantareira since 1901; of these, nine have not been sighted there for the last two decades. The number of bird species endemic to the Atlantic Forest is high (80), and seven of its species are globally threatened. According to multivariate analyses the species diversity at Serra da Cantareira is similar to other regions of the Atlantic Forest, such as Carlos Botelho and Intervales state parks, where the vegetation is also ombrophilous dense forest. We discuss local changes in the avifaunal composition over the last decades and suggest the incorporation of large forest remnants to the Cantareira State Park to mitigate the impact of the northern section of Rodoanel Mário Covas, a highway (SP-21) that will soon be operational and will negatively impact the biodiversity of Serra da Cantareira.

KEY WORDS. Atlantic Forest, bird conservation, hierarchical cluster analysis, principal coordinate analysis, Serra da Cantareira.

INTRODUCTION

The Atlantic Forest (AF) is the second largest rainforest in South America and has a rich and diverse avifauna (900 species, 24% of which are endemic to the AF; Lima 2013). New bird species are still being discovered in this forest, even near large urban areas (Buzzetti et al. 2013), demonstrating that the local avifauna is not completely known, and reinforcing the importance of bird surveys there. Despite the high levels of species richness and endemism, only ~12% of the AF forest remains, and as a result, a large number of its component taxa are at imminent risk of extinction (Marini and Garcia 2005, Ribeiro et al. 2009). In addition to the importance of the AF for

bird conservation, its forest remnants provide several ecosystem services for human populations, such as guaranteeing water supply near major urban areas (Dean 1995).

After 1893, due to a water collapse in the city São Paulo, the government started to expropriate farms at Serra da Cantareira for reforestation, to ensure the recovery of the streams and river that supply the city with water (Vilar 2007). Considered the central piece in the São Paulo City Green Belt Biosphere Reserve by UNESCO and an Important Bird Area by BirdLife International (IBA SP03), Serra da Cantareira still provides water and other ecosystem services to the city, and houses a significant portion of the AF biodiversity, including several threatened and endemic species (Bencke et al. 2006, Whately and Cunha




Figure 1. Serra da Cantareira locations surveyed in this study. In the urban slope are Cantareira State Park administrative zones and their trails: AC = Águas Claras; PG = Pedra Grande; Eng = Engordador; Cab = Cabuçu; EL = Estrada do Lenhador; Alberto Löfgren State Park (ALSP) and A2. In the inner slope are A1, A3, A4, A5 and SC = Sítio Capuavinha private reserve.

2007, Ayres 2008). Serra da Cantareira stretches for more than 10,000 ha in the Metropolitan Area of the city of São Paulo, the most populated and urbanized region in South America, and currently comprises the largest native urban rainforest in the world (Ayres 2008). The main impacts to the biodiversity of Serra da Cantareira are urban expansion and large infrastructure projects in nearby towns (Bencke et al. 2006, Fig. 1).

Although Serra da Cantareira is continuous with one of the largest cities in the world and near several research institutions and universities, few ornithological studies have been performed there. Museum collectors were the first to survey these mountains in the end of the XIX century (see Pinto 1938, 1944, Paynter and Traylor 1991). Graham (1992) published the only comprehensive bird inventory of the region, more than 20 years ago. Other than museum specimens and Graham's article, there have been few ornithological studies and bird surveys. They range from comparison of communities in different vegetation types (Graham 1991, Antunes et al. 2009; frugivory (Fonseca and Antunes 2007, Ikuta and Martins 2013); new species records (Antunes et al. 2008a), plumage anomaly (Cavarzere and Tonetti 2015); nest description (V.R. Tonetti et al. unpublished data),

studies on species ecology (Tonetti and Pizo 2016), technical reports (Antunes and Eston 2009, 2012, DERSA 2010) and records provided by birdwatchers (WikiAves 2016, Xeno-Canto 2016).

In this article, we present an updated compilation of all bird species recorded for Serra da Cantareira, including our personal records. We compare the avifauna between Serra da Cantareira and other regions of AF using multivariate analysis, discuss changes in avifaunal composition during the last decades, and provide additional information on some threatened and/or uncommon species. Lastly, we focus on specific localities and measures for bird conservation in our study area.

MATERIAL AND METHODS

Serra da Cantareira ranges from 750 to 1,250 m asl. The vegetation there is predominantly montane ombrophilous dense forest. The climate is mesothermal and humid, with rainy summers and dry winters (CWA Köppen), and annual temperature and rainfall averaging 20 °C and 1,500 mm, respectively (Bencke et al. 2006). The slope facing the urban area comprises the most preserved area, with continuous forests, which is mostly



protected by Cantareira (7,900 ha; hereafter CSP) and Alberto Löfgren state parks (187 ha; ALSP). CSP is divided into four administrative zones: Águas Claras, Cabuçu, Engordador and Pedra Grande (Fig. 1). Patches in the inner slope have variable sizes (~1 to 100 ha) and are at different stages of regeneration. In this study we considered bird records from CSP, ALSP, neighboring patches in the inner slope, including the 20-ha private reserve Sítio Capuavinha (Lemos 2014) and records that indicated only "Serra da Cantareira" as a location, without any further detail.

We obtained species records over the last decade during systematic and unsystematic bird surveys. In unsystematic surveys, samplings consisted of recording all aurally or visually detected birds at unlimited distance, while walking on trails (Ribon 2010). Surveys were performed during the morning in different seasons and were mostly concentrated on the available trails of the four CSP administrative zones and a dirt road named "estrada do lenhador", at the park's boundaries (Fig. 1). Birds were recorded with the aid of binoculars and song recordings and photographs were occasionally taken during surveys.

VRT performed a systematic survey using point-counts. In this study, 100 fixed-points were established in the four CSP administrative zones and visits lasted 10 minutes during the morning. Each point was visited six times on different days between May and December 2014, resulting in a total of 600 samples, which include only birds detected within a 50-m radius (Vielliard et al. 2010, Tonetti and Pizo 2016). MAR, along with other researchers (see names in acknowledgements), performed another systematic study using line transects (80 h of sampling effort) and mist-nets (2,600 net-hours) in five different areas (A1 to A5; Fig. 1). These surveys were conducted during the morning and late afternoon from August 12th 2009 to April 29th 2010. Some netted specimens were collected and deposited in the MZUSP collection (details in DERSA 2010).

In order to compile our data, we searched for museum specimens collected from "Cantareira State Park", "Parque Estadual da Cantareira", "Cantareira Mountains", "Serra da Cantareira", "Cantareira", "Alberto Löfgren State Park", "Parque Estadual Alberto Löfgren", "Horto Florestal", "Cuca, Horto Florestal" and "Parque Florestal". We surveyed the collections of the following museums: Museum of Zoology of the University of São Paulo (MZUSP, Brazil), Museum of Zoology of the Campinas University (MUNICAMP, Brazil), and the Field Museum of Natural History (FMNH, USA). The localities cited above correspond to CSP, ALSP or not specified sites at Serra da Cantareira (Paynter and Traylor 1991). Using the same terms as outlined above, we searched for indexed articles potentially containing bird records in the citation databases Web of Science, Google Scholar and SciELO. Additionally, we used Google to search for non-indexed articles and "gray" literature, such as technical reports and theses. We also compiled records from the WikiAves (www.wikiaves.com; WA) and Xeno-canto (www.xeno-canto.org; XC) databases. All searches were performed until June 1st 2016. We discarded erroneous or doubtful records – i.e. misidentified pictures and

song records, as well as species pending confirmation of their occurrence in our study area according to the literature. We followed the nomenclature of the CBRO (Brazilian Ornithological Records Committee) (Piacentini et al. 2015).

We used Mountford's dissimilarity index (Wolda 1981) to perform a hierarchical cluster analysis (HCA) and a Principal Coordinate Analysis (PCoA) to compare the bird communities from Serra da Cantareira with other AF regions. After calculating the Mountford's index using presence/absence data with the *vegdist* function in the R package *vegan* (Oksanen et al. 2016), we performed the HCA using the Ward's minimum variance criterion as the objective function, the results of which define which clusters merge at each step (Ward Jr 1963). The HCA and the PCoA were performed using the *hclust* and *ordiplot* functions (Oksanen et al. 2016, R Core Team 2016). We also calculated the Jaccard dissimilarity index using the function *vegdist* (Oksanen et al. 2016). The Jaccard coefficient ranges from 0 (identical pairwise) to 1 (totally different pairwise) and it is a robust index to ascertain similarity among groups (Krebs 1999).

We compared bird communities between Serra da Cantareira and the following AF regions that also have dense ombrophilous forest: Boraceia Biological Station (Cavarzere et al. 2010), Carlos Botelho State Park (Antunes 2013), Intervales State Park (Antunes et al. 2008b), Morro Grande Reserve (Develey and Martensen 2006), Municipality of Ilha Comprida (Gussoni 2010), Municipality of Ubatuba (Simpson et al. 2012), and in Serra da Mantiqueira, where the vegetation is predominantly mixed ombrophilous forest: Campos do Jordão (composed by the Mananciais de Campos do Jordão and Campos do Jordão State Parks; Willis and Oniki 1981) and Itatiaia National Park (Barreto et al. 2013), as well as regions of semi-deciduous forest: Barreiro Rico Ecological Station (Antunes 2005), Caetetus Ecological Station (Tabanez et al. 2005, Cavarzere et al. 2009), Mata dos Godoy State Park (Anjos 2001), Morro do Diabo State Park (Uezu and Metzger 2016), Porto Ferreira (composed by Porto Ferreira and Vassununga State Parks, Uezu and Gaban-Lima 2003) and Rio Claro Farm (Donatelli et al. 2004). In addition to the references cited above, to complement lists of birds from those regions, we also used records from WikiAves and Xeno-Canto databases (WikiAves 2016, Xeno-Canto 2016). Most of those regions are or incorporate large conservation areas (reserves) where bird richness is high. Additionally, whenever possible, we also compiled records from patches surrounding the parks and reserves, to better characterize the avifauna, in the same manner that we did for Serra da Cantareira. These regions are ~230 km distant from our study area (50 km the nearest and 600 km the farthest; Fig. 2). For comparisons, we ignored seabirds records for Boracéia Biological Station and municipalities of Ilha Comprida and Ubatuba (e.g. the Magnificent Frigatebird *Fregata magnificens* [Mathews, 1914]) as well as exotic species, such as the House Sparrow *Passer domesticus* (Linnaeus, 1758), and birds from other regions in Brazil that escaped or were intentionally released from captivity (e.g. the Yellow-rumped Cacique *Cacicus cela* [Linnaeus, 1758]).




Figure 2. Dendrogram with Atlantic Forest regions based on their bird community using Hierarchical Cluster Analysis (Mountford's distance coefficient; a) and the Principal Coordinate Analysis of these areas (b). 1 = Morro do Diabo State Park; 2 = Mata dos Godoy State Park; 3 = Caetetus Ecological Station; 4 = Rio Claro Farm; 5 = Intervales State Park; 6 = Barreiro Rico Ecological Station; 7 = Carlos Botelho State Park; 8 = Ilha Comprida; 9 = Morro Grande Reserve; 10 = Porto Ferreira (Porto Ferreira and Vassununga State Parks); 11 = Serra da Cantareira; 12 = Boracéia Biological Station; 13 = Campos do Jordão (composed by Mananciais de Campos do Jordão and Campos do Jordão State Parks); 14 = Ubatuba; 15 = Itatiaia National Park.

RESULTS

Overall, 326 species were found at Serra da Cantareira, including seven exotic species, e.g. the Rock Pigeon *Columba livia* (Gmelin, 1789), or species that may have escaped from captivity, such as the Red-cowled Cardinal *Paroaria dominicana* (Linnaeus, 1758). For nine species with previous records for Serra da Cantareira there have been no further records for the last two decades, and 55 species records are not adequately documented (through either museum specimen, photograph or song record; Appendix 1). Additionally, nine questionable literature records of bird species, for instance the Vinaceous Parrot *Amazona vinacea* (Kuhl, 1820), were not included in this report (Appendix 2). Non-passerines accounted for 138 species in 32 families and passerines are represented by 188 species in 34 families (Piacentini et al. 2015). Tyrannidae was the most representative family (35

species), followed by Thraupidae (33) and Furnariidae (18). AF endemics accounted for 24.5% of the records according to Lima's (2013) classification and 227 are forest birds (Parker et al. 1996). Seven species recorded in our study area are globally threatened (BirdLife 2016); five are threatened in Brazil (MMA 2014) and 17 in the state of São Paulo (São Paulo 2014). The ornithological knowledge of Serra da Cantareira has increased significantly since Graham's study. Only 71 birds were reported before 1992, all represented by museums specimens (Fig. 3). Records exclusively from the authors of this study contributed to add four species (Appendix 1). Species in the urban slope accounted for most records (Table 1).

HCA showed two major groups of areas, separated according to their bird communities. One is in a macro-region where the vegetation is predominantly ombrophilous dense or ombrophilous mixed forest near the coast, and comprises the




Figure 3. Number of species reported by different sources during different periods in Serra da Cantareira. Museums: specimens deposited in ornithological collections between 1901 and 1992; Graham: Graham 1992; Authors: species recorded by the authors during unsystematic surveys; WA: WikiAves 2016; XC: Xeno-Canto 2016; Recent literature: Figueiredo and Loo 2000, Develey and Endrigo 2004, Fonseca and Antunes 2007, Antunes et al. 2008a, 2008b, 2009, Antunes and Eston 2009, 2012, Minns et al. 2009, Dersa 2010, Ikuta and Martins 2013, Lemos 2014, Cavarzere and Tonetti 2015. All recent encompasses records obtained in recent literature cited above as well as specimens deposited in museums after the work performed by Graham (1992) and species recorded by the authors.

Table 1. Number of species recorded in the different slopes of Serra da Cantareira. Urban slope corresponds to species recorded in Cantareira and Alberto Löfgren State Parks. Inner slope corresponds to species recorded in the slope faced to inner Atlantic forest in A1, A3, A4, A5 and the private reserve Sítio Capuavinha. In this table we did not take into account species which the location pointed only to Serra da Cantareira.

Locality	Number of recorded species
Urban slope	298
CSP	293
ALSP	207
Inner slope	222
A1, A3, A4 and A5	202
Sítio Capuavinha reserve	98

following areas: Boracéia Biological Station, Campos do Jordão, Serra da Cantareira, Carlos Botelho and Intervales State Parks, Ilha Comprida, Itatiaia National Park, Morro Grande Reserve and Ubatuba. The other group, encompassing more inland areas,

has lower rainfall rates with seasonal semi-deciduous forests and includes: Barreiro Rico Biological Station, Caetetus Ecological Station, Mata dos Godoy State Park, Morro do Diabo State Park, Porto Ferreira and Rio Claro Farm (Fig. 2). The bird community of Serra da Cantareira is more similar to a clade composed of Boracéia Biological Station, Carlos Botelho and Intervales state parks and Morro Grande Reserve (Fig. 1a) according to HCA. According to the Jaccard index the bird community of Serra da Cantareira is more similar to Carlos Botelho State Park, while Pontal do Parapanapanema has a more distinct avifauna (Table 2). Although the ombrophilous mixed forests, represented here by Campos do Jordão and Itatiaia National Park, are more related to ombrophilous dense forests according to HCA, they form a separated clade within this major group (composed by areas 13 and 15, Fig. 2a). The PCoA also showed two regions of ombrophilous mixed forests that are more related to ombrophilous dense forests in the first dimension, while in the coordinate 2 they were more distant from other ombrophilous dense forest areas (Fig. 2b).

Table 2. Areas compared with Serra da Cantareira in multivariate analysis using presence/absence bird species data. spp. = number of species reported; Distance (km) = distance in kilometers from our study area; Jaccard = Jaccard dissimilarity index, the lower the number, the greater the similarity in avifauna composition with Serra da Cantareira. Areas with avifauna most (Carlos Botelho State Park) and least (Morro do Diabo State Park) similar to Serra da Cantareira are indicated in bold.

Area	Extension (ha)	spp.	Distance (km)	Jaccard
Barreiro Rico Ecological Station	300	198	180	0.557
Boracéia Biological Station	96	313	75	0.392
Caetetus Ecological Station	2,180	290	330	0.491
Carlos Botelho State Park	38,000	333	150	0.354
Campos do Jordão	9,000	233	130	0.514
Ilha Comprida	19,200	257	190	0.543
Intervales State Park	42,000	421	190	0.377
Itatiaia National Park	24,000	404	220	0.393
Mata dos Godoy State Park	680	268	470	0.474
Morro Grande Reserve	10,000	194	50	0.498
Morro do Diabo State Park	34,000	215	600	0.616
Porto Ferreira	2,700	240	200	0.545
Rio Claro Farm	4,000	219	260	0.566
Ubatuba	71,078,300	377	140	0.407

DISCUSSION

Our results are consistent with other ornithological compilations conducted at Boracéia Biological Station, which is an AF reserve located at Serra do Mar, 75 km distant from Serra da Cantareira, and where the vegetation is also dense mountain rain forest (Fig. 2). Cavarzere et al. (2010) performed transect counts, mist netting and unsystematic surveys at the station, where they recorded 242 bird species. After accounting for



museums specimens, literature records, and records provided by birdwatchers, 323 species were reported for that reserve, 30% of which are endemic to the AF (Cavarzere et al. 2010, Lima 2013). This suggests that ornithological reviews provide a more comprehensive knowledge on the actual bird diversity of a given region than localized surveys, even when using different methodologies and intensive sampling effort.

Some records of species for Serra da Cantareira are based on only one source of information (records from the authors, those provided by birdwatchers, reported in the literature or represented by museums specimens; Fig. 3), e.g. the Gray-bellied Hawk *Accipiter poliogaster* (Temminck, 1824). This bird was only recently spotted by the authors and had not been previously recorded there. Despite its broad-range distribution (north Colombia to south Brazil, and more recently found in Costa Rica and Panama), this species is considered uncommon and is sparsely found throughout its range (Bierregard et al. 2016). VRT first recorded a juvenile at Cabuçu (a CSP administrative zone; Fig. 1) on May 5, 2014. On that day, the birds' vocalization was recorded.

The high number of species recorded by birdwatchers (76% of all species compiled in this study, ten of which reported only by them; Appendix 1) highlights the importance of public websites where digital vouchers can be deposited, such as WikiAves (<http://www.wikiaves.com>) and Xeno-Canto (<http://www.xeno-canto.org>). Among the records provided only by these databases is the Shrike-like Cotinga *Laniisoma elegans* (Thunberg, 1823) (WA1076770). This frugivorous bird, found mostly in old-growth forests, was photographed in a forest edge at the Engordador administrative CSP zone. Birds recorded only in the literature accounted for 13 species, such as the Rufous-tailed Attila *Attila phoenicurus* Pelzeln, 1868, song-recorded at Cabuçu (Minns et al. 2009). The Rufous-tailed Attila breeds in southeast Brazil during the austral winter, and is found in greater numbers in Serra do Mar (Walther 2016), being uncommon in our study area (pers. obs.).

Although museum records until 1992 did not account for 80% of the species in Serra da Cantareira, some relevant birds were reported only until that year. The most noticeable is the Purple-winged Ground-Dove *Claravis geoffroyi* (Temminck, 1811) collected in 1937 at a CSP site named "Cuca" (MZUSP 17040; Fig. 1). It was once a fairly common and widely distributed species (occurring from Bahia [Brazil] south through eastern Brazil to northern Argentina and eastern Paraguay). During the last three decades, however, there have been only a few non-documented records of this globally "Critically Endangered" AF endemic species (MMA 2014, BirdLife 2016). *Claravis geoffroyi* feeds on bamboo seeds (*Guadua* sp.), and most likely travels far in search for flowering events (Sick 1997, Areta et al. 2009). *Guadua* spp. have synchronous masting events in cycles of ~30 y, attracting birds that eat their seeds, such as the Buffy-fronted Seedeater *Sporophila frontalis* (Verreaux, 1869) and the Temminck's Seedeater *S. falcirostris* (Temminck, 1820), both registered at Serra da Cantareira and globally threatened (Areta et al. 2009, BirdLife 2016). As *Guadua* sp. bamboo tickets are common in several parts

of Serra da Cantareira (Bencke et al. 2006; pers. obs.), we suggest that those searching for the Purple-winged Ground-Dove should try to locate flowering events of that bamboo.

Due to severe forest loss and human intervention at Serra da Cantareira, it is possible that local extinctions have occurred, but they are difficult to document. For example, Cavarzere et al. (submitted) recorded the Squamate Antbird *Myrmotherus squamosus* (Pelzeln, 1868) after a 190-y time span from the last confirmed record at the Ipanema National Forest, an AF reserve 100 km from our study area. This is a good example of imperfect detection and insufficient sampling effort (Mackenzie et al. 2003). The Spot-billed Toucanet *Selenidera maculirostris* (Lichtenstein, 1823) is an example of a species recently recorded at Serra da Cantareira after 45 y without records. Six specimens were collected in 1965 (MZUSP 60592 to 60597), and after it, the only confirmed record was provided by MAR and his team in A5 (Fig. 1).

More important than accounting for local bird extinctions *per se* is identifying functional extinctions and their impact on the ecosystem. Habitat disturbance is one of the main drivers of functional extinctions of sensitive guilds, such as understory insectivorous and large-bodied frugivorous (Martensen et al. 2012, Galetti et al. 2013, Morante-Filho et al. 2015). It is very possible that, of the currently rarest species in our study area – e.g. the Saffron Toucanet *Pteroglossus bailloni* (Vieillot, 1819), a large-bodied frugivorous, and the Speckle-breasted Antpitta *Hylomanes nattereri* (Pinto, 1937), an understory insectivorous – were more abundant in the past, given their presence in museums collections. Declines in the populations of these birds have been recorded in other AF regions due to habitat disturbances (Guix et al. 2000, Anjos 2006), and since they participate in key-ecological processes (e.g. seed dispersal), the consequences of their functional extinctions should be better investigated.

In contrast with the decreasing populations of some species in response to habitat disturbances, species that benefit from altered environments are expected to expand their ranges and increase their abundance. Graham (1992), using point-counts (37 hours of sampling effort) between December 11th 1985 and March 29th 1986, did not record any individual of the White-eyed Parakeet *Psittacara leucophthalmus* (Statius Muller, 1776) within the CSP, although this species has been frequently recorded in our study area over the last decade. The White-eyed Parakeet benefits from deforestation and occupies forest edges and areas of open vegetation, and its presence in AF is an indication of environmental degradation (Sick 1997). Increases in the numbers of generalist bird species, together with decreases in the numbers of sensitive species, have also been reported for other large and protected AF remnants after a 30-y time span – at the Caetetus Ecological Station (Cavarzere et al. 2012; 2,800 ha, Fig. 2).

Biogeographic affinities

Similarly to a study on the plants of Serra da Cantareira, our results indicate that the local avifauna is associated more strongly



with dense ombrophilous forest areas, to a lesser extent to mixed ombrophilous forest and to an even lesser extent to seasonal semi-deciduous forests (Salis et al. 1995; Fig. 2a, Table 2). The Black-backed Tanager *Tangara peruviana* (Desmarest, 1806) and the Olive-green Tanager *Orthogonyx chloricterus* (Vieillot, 1819) are typical of the dense ombrophilous forest species of Serra da Cantareira that occur predominantly near the coast. However, birds commonly found in seasonal semi-deciduous forests, such as the Southern Antpitta *Corythopis delalandi* (Lesson, 1830), were also recorded. This understory insectivorous species has been song-recorded only once by the MAR team at the administrative zone of Pedra Grande at CSP. In order to maintain the genetic flow and evolutionary processes between these three types of vegetation, the establishment of forest corridors uniting them is crucial. Forest patches are important in facilitating bird dispersal over landscapes and can help to connect larger AF remnants (Ribeiro et al. 2009).

Conservation

In addition providing information on the bird community of Serra da Cantareira, our results reinforce the relevance of the area for birds. Bencke et al. (2006) estimated an avifauna richness of 250 species for the region and reported 65 AF endemics and four globally threatened species, less than the numbers provided by us (326, 80 and seven respectively). Although located at an urban matrix and composed mostly of secondary forest, Serra da Cantareira accounted for 17% of all species recorded in Brazil (Piacentini et al. 2015) and 36.7% in AF (Lima 2013). Moreover, it is a stronghold for populations of some species that are suffering a sharp decline in several other AF regions, for instance the Solitary Tinamou *Tinamus solitarius* (Vieillot, 1819), currently found mostly in large forest remnants such as those in Serra do Mar (BirdLife 2016) and also an abundant bird in our study area (Bencke et al. 2006, pers. obs.), as well as the Southern Bristle-tyrant *Phylloscartes eximus* (Temminck, 1822), a passerine that was once broadly distributed but now is found only in a few places (Silveira 2009, Tonetti and Pizo 2016).

Despite differences in sampling effort, we found that the avifauna composition of CSP and that of the inner slope patches are similar (Appendix 1), corroborating the hypothesis that even fragmented landscapes with high levels of forest cover can maintain similar species richness to areas of continuous forest (Martensen et al. 2012, Morante-Filho et al. 2015). This supports the idea that forest patches surrounding Serra da Cantareira should be protected, since they can act as corridors between our study area and other large AF remnants, such as those in Serra da Mantiqueira and Serra do Mar (Fig. 2). Since seasonal semi-deciduous forests have been more extensively devastated than ombrophilous and mixed forests, there are only a few seasonal semi-deciduous forest remnants that are larger than 500 ha with weakly connected patches (Ribeiro et al. 2009), which means that genetic flow between Cantareira birds and those that inhabit the inner AF can be seriously compromised.

Similarly to Serra da Cantareira, other large (15,700 ha) urban forests, in the city Rio de Janeiro City, the Tijuca National Forest and the Pedra Branca State Park, have high bird richness (325 species; Lepage 2016), and are also considered Important Bird Areas (IBA RJ07, Bencke et al. 2006), with seven globally threatened species. This highlights the importance of such reserves, which despite the high levels of anthropic pressure, harbor a significant portion of our biodiversity. As in our study area, the main threats to the Tijuca National Forest are urban sprawl, and the impacts of large infrastructure projects in the surrounding cities (Bencke et al. 2006).

When it comes to infrastructure, roads cause one of the largest disturbances in urban forests. Their intense noise can cause changes in the foraging behavior of birds, hinder intra and inter-specific communication, and cause chronic stress. The animals, in turn, become more prone to diseases and their reproductive success is reduced (Ortega 2012). There is one highway (Fernão Dias, BR-116) and tree paved roads with intense traffic crossing CSP (da Silva et al. 2009). In addition to this, the northern section of Rodoanel Mário Covas (SP-21) will be operating soon (DERSA 2010; Fig. 1). This new highway is being built close to the boundaries of CSP and ALSP, crossing the first in some sections via tunnels, and will represent a new threat to the biodiversity of Serra da Cantareira (Ayres 2008, da Silva et al. 2009, DERSA 2010; Fig. 1). As a strategy to mitigate the damage caused by those roads, we suggest the incorporation of forest remnants to the CSP, especially those large patches in the west portion of the Park, and near Cuca (Fig. 1).

ACKNOWLEDGEMENTS

We thank the staff of Cantareira State Park for logistic assistance; the scientific technical committee (COTEC) which authorized VRT to work on the Park (proc. 260108-007.094/2013); Instituto Chico Mendes de Conservação da Biodiversidade (ICMBio) for collecting permits; the Brazilian National Council for Scientific and Technological Development which provided a scholarship to VRT (proc. 130279/2013-7) and for grants and financial support to LFS; people who contribute in the maintenance as well as providing their personal records to WikiAves and Xeno-Canto databases; MAR team: Aline Correa, Fernanda Alves, Fernanda Bocalini, Giuliana Althman, Glaucia Del Rio, Leo Signorini, Rafael Marcondes, Thiago Vernaschi and Wagner Cavarzere. We are also indebted with Tatiana Pongiluppi. Fernanda Alves helped with the English. Maria de Fátima Rodrigues funded part of this research.

LITERATURE CITED

- Anjos L (2001) Bird communities in five Atlantic forest fragments in southern Brazil. *Ornitología Neotropical* 12: 11–27.
- Anjos L (2006) Bird species sensitivity in a fragmented landscape of the Atlantic forest in southern Brazil. *Biotropica* 38: 229–234. <https://doi.org/10.1111/j.1744-7429.2006.00122.x>

- Antunes AZ (2005) Alterações na composição da comunidade de aves ao longo do tempo em um fragmento florestal no sudeste do Brasil. *Ararajuba* 13: 47–61.
- Antunes AZ (2013) Aves do Parque Estadual Carlos Botelho. *Biota Neotropica* 13: 124–140. <https://doi.org/10.1590/S1676-06032013000200012>
- Antunes AZ, Eston MR (2009) Anexo 10: espécies de aves registradas no PEC. In: Leonel C, Déscio F, Mazzei K, Leite MO, Barros MIA (Eds). *Parque Estadual da Cantareira: plano de manejo*. São Paulo, Secretaria de Estado de Meio Ambiente, Instituto Florestal, Fundação Florestal, 1–10.
- Antunes AZ, dos Santos AMR, Eston MR (2008a) Registros relevantes de aves para a grande São Paulo obtidos no Parque Estadual da Cantareira (Nota Científica). *Revista do Instituto Florestal* 20: 221–224.
- Antunes AZ, Menezes GV, Eston MR (2008b) Avifauna. In: Furlan SA, Leite SA (Eds) *Parque Estadual de Intervales: plano de manejo*. São Paulo, Secretaria de Estado de Meio Ambiente, Instituto Florestal, Fundação Florestal, 122–138.
- Antunes AZ, Wyrgun B, Eston MR (2009) Composição das comunidades de aves em duas florestas secundárias contíguas no sudeste do Brasil. *Revista do Instituto Florestal* 21: 93–106.
- Antunes AZ, Eston MR (2012) Anexo 8: espécies de aves registradas no PEAL. In: Arromba AL, Leonel C, Santiago CM, Mazzei K, Bucci LA, Nalon MA, Barros MIA (Eds) *Parque Estadual Alberto Löfgren: plano de manejo*. Secretaria de Estado de Meio Ambiente, Instituto Florestal, Fundação Florestal, São Paulo, 1–10.
- Areta JI, Bodrati A, Cockle K (2009) Specialization on guadua bamboo seeds by three bird species in the Atlantic forest of Argentina. *Biotropica* 41: 66–73. <https://doi.org/10.1111/j.1744-7429.2008.00458.x>
- Ayres ACM (2008) O Ciclo da Caapora: A RMSP e o Parque da Cantareira. Annablume, São Paulo, 120 pp.
- Barreto CG, Campos JB, Roberto DM, Schwarzstein NT, Alves GSG, Coelho WC (2013) Plano de manejo: Parque Nacional do Itatiaia. Instituto Chico Mendes de Conservação da Biodiversidade, Ministério do Meio Ambiente, Brasília, 215 pp.
- Bencke GA, Maurício GN, Develey PF, Goerck JM (2006) Áreas importantes para a conservação das aves no Brasil: parte 1 – estados do domínio da Mata Atlântica. SAVE Brasil, São Paulo, 494 pp.
- Bierregard RO, Christie DA, Kirwan G (2016) Grey-bellied Goshawk (*Accipiter poliogaster*). In: del Hoyo J, Elliot A, Sargatal J, Christie DA, de Juana E (Eds) *Handbook of the birds of the world alive*. Available online at: <http://www.hbw.com/node/53041> [Accessed: 27/05/2016]
- BirdLife International (2016) IUCN Red List for Birds. <http://www.birdlife.org> [Accessed: 19/03/2016]
- Buzzetti DRC, Belmonte-Lopes R, Reinert BL, Silveira LF, Bornschein MR (2013) A new species of Formicivora Swainson, 1824 (Thamnophilidae) from the state of São Paulo, Brazil. *Revista Brasileira de Ornitologia* 21: 269–291.
- Cavarzere V, Marcondes RS, Moraes GP, Donatelli RJ (2012) Comparação quantitativa da comunidade de aves de um fragmento de floresta semidecidual do interior do Estado de São Paulo em intervalo de 30 anos. *Iheringia* 102: 384–393. <https://doi.org/10.1590/S0073-47212012000400004>
- Cavarzere V, Moraes GP, Donatelli RJ (2009) Avifauna da Estação Ecológica dos Caetetus. *Papéis Avulsos de Zoologia* 49: 477–485.
- Cavarzere V, Moraes GP, Silveira LF (2010) Boracéia Biological Station: an ornithological review. *Papéis Avulsos de Zoologia* 50: 189–201. <https://doi.org/10.1590/S0031-10492010001300001>
- Cavarzere V, Tonetti VR (2015) Complete leucism in the Fawn-breasted Tanager *Pipraeidea melanonota* (Aves: Thraupidae). *Boletim do Museu de Biologia Prof. Mello Leitão* 37: 233–240.
- da Silva DA, Pavão M, Fernandes A, Zuquim ML, Miaciro K, Gasparotto TC, Veiga MF (2009) Avaliação do meio antrópico. In: Leonel C, Déscio F, Mazzei K, Leite MO, Barros MIA (Eds) *Parque Estadual da Cantareira: plano de manejo*. Secretaria de Estado de Meio Ambiente, Instituto Florestal, Fundação Florestal, São Paulo, 145–222.
- Dean W (1995) *A ferro e fogo: a história de devastação da Mata Atlântica brasileira*. Companhia das letras, São Paulo, 484 pp.
- DERSA (2010) Programa Rodoanel Mario Covas, Trecho Norte: Estudo de Impacto Ambiental. Volume IV. Desenvolvimento Rodoviário S/A, Consórcio JPGP-PRIME, São Paulo, 451 pp.
- Develey PF, Endrigo E (2004) *Guia de Campo – Aves da Grande São Paulo*. Editora Aves e Fotos, São Paulo, 295 pp.
- Develey PF, Martensen AC (2006) As aves da Reserva do Morro Grande (Cotia, SP). *Biota Neotropica* 6: 1–16. <https://doi.org/10.1590/S1676-06032006000200008>
- Donatelli RJ, da Costa TVV, Ferreira CD (2004) Dinâmica da avifauna em um fragmento de mata na Fazenda Rio Claro, Lençóis Paulista, São Paulo, Brasil. *Revista Brasileira de Zoologia* 21: 97–114. <https://doi.org/10.1590/S0101-81752004000100017>
- Figueiredo LFA, Loo VK (2000) Lista das aves do município de São Paulo. *Boletim do CEO* 14: 15–35.
- Fonseca FY, Antunes AZ (2007) Frugivoria e predação de sementes por aves no Parque Estadual Alberto Löfgren, São Paulo, SP. *Revista do Instituto Florestal* 19: 81–91.
- Galetti M, Guevara R, Côrtes MC, Fadini R, Von Matter S, Leite AB, Labeca F, Ribeiro T, Carvalho CS, Collevatti RG, Pires MM, Guimarães PR, Brancalion PH, Ribeiro MC, Jordano P (2013) Functional Extinction of Birds Drives Rapid Evolutionary Changes in Seed Size. *Science* 340: 1086–1090. <https://doi.org/10.1126/science.1233774>
- Graham DJ (1991) The avifauna and the vegetation structure of a mature araucaria plantation in São Paulo, Brazil. *IF, Série Registros* 6: 1–79.
- Graham DJ (1992) The Avifauna of the Serra da Cantareira, São Paulo, Brazil: A preliminary survey. *IF, Série Registros* 10: 1–56.
- Guix JC, Martín M, Hernández A, Souza FL (2000) Conservation status of the Saffron Toucanet (*Baillonius bailloni*, Ramphastidae): a new case of population isolation and depletion in South America. *Grupo de Estudos Ecológicos, Série Documentos* 6: 10–25.



- Gussoni COA (2010) Novas informações sobre a história natural da maria-da-restinga (*Phylloscartes kronei*) (Aves, Tyrannidae). Master dissertation, Universidade Estadual Paulista Júlio de Mesquita Filho, Rio Claro. Available online at: https://repositorio.unesp.br/bitstream/handle/11449/99487/gussoni_coa_me_rcla.pdf?sequence=1&isAllowed=y [Accessed: 19/03/2016]
- Ikuta KG, Martins FC (2013) Interação entre aves frugívoras e plantas no Parque Estadual da Cantareira, estado de São Paulo. *Atualidades Ornitológicas* 172: 33–36.
- Krebs CJ (1999) Ecological Methodology. Addison-Wesley Educational Publisher, Menlo Park, 642 pp.
- Lemos RF (2014) Avifauna da Reserva Particular do Patrimônio Natural (RPPN) Sítio Capuavinha, município de Mairiporã, São Paulo. *Atualidades Ornitológicas* 179: 38–45.
- Lepage D (2016) Parque Nacional da Tijuca, Rio de Janeiro. Avibase – Bird Checklists of the World. <http://avibase.bsc-eoc.org/checklist.jsp?region=BRRj01> [Accessed: 19/03/2016]
- Lima LM (2013) Birds of the Atlantic Forest: richness, composition, status, endemism, and conservation. Master dissertation, Universidade de São Paulo, São Paulo. <http://www.teses.usp.br/teses/disponiveis/41/41133/tde-17042014-091547/pt-br.php> [Accessed: 19/03/2016]
- Mackenzie DL, Nichols JD, Hines JE, Knutson MG, Franklin AB (2003) Estimating site occupancy, colonization, and local extinction when a species is detected imperfectly. *Ecology* 84: 2200–2207. <https://doi.org/10.1890/02-3090>
- Marini MA, Garcia FI (2005) Bird conservation in Brazil. *Conservation Biology* 19: 665–671. <https://doi.org/10.1111/j.1523-1739.2005.00706.x>
- Martensen AC, Ribeiro MC, Banks-Leite C, Prado PI, Metzger JP (2012) Associations of forest cover, fragment area, and connectivity with neotropical understory bird species richness and abundance. *Conservation Biology* 26: 1100–1111. <https://doi.org/10.1111/j.1523-1739.2012.01940.x>
- Minns JC, Buzzetti DRC, Albano CG, Whittaker A, Gross AE, Parolini R (2009) Birds of Brazil/Aves do Brasil. Aves Brasiliis Editora, Vinhedo, DVD-ROOM.
- MMA (2014) Lista de espécies ameaçadas. Ministério do Meio Ambiente. <http://www.icmbio.gov.br> [Accessed: 18/03/2016]
- Morante-Filho JC, Faria D, Mariano-Neto E, Rhodes JR (2015) Birds in anthropogenic landscapes: the responses of ecological groups to forest loss in the Brazilian Atlantic forest. *PLoS ONE* 10: e0128923. <https://doi.org/10.1371/journal.pone.0128923>
- Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'Hara RB, Simpson GL, Solymos P, Stevens MHH, Wagner H (2016) vegan: Community Ecology Package. <https://CRAN.R-project.org/package=vegan> [Accessed: 18/03/2016]
- Ortega CP (2012) Effects of noise pollution on birds: a brief review of our knowledge. *Ornithological Monographs* 74: 6–22. <https://doi.org/10.1525/om.2012.74.1.6>
- Parker TAI, Stotz DF, Fitzpatrick JW (1996) Ecological and Distributional Databases. In: Stotz DF, Fitzpatrick JW, Parker TAI, Moskovits DA (Eds) *Neotropical Birds: Ecology and Conservation*. The University of Chicago Press, Chicago, 133–293.
- Paynter RA, Traylor MA (1991) *Ornithological Gazzeteer of Brazil*. Cambridge, Harvard University, 788 pp.
- Piacentini VQ, Aleixo A, Agne CE, Maurício GN, Pacheco JF, Bravo GA, Brito GRR, Naka LN, Olmos F, Posso S, Silveira LF, Betini GS, Carrano E, Franz I, Lees AC, Lima LM, Pioli D, Schunck F, Amaral FR, Bencke GA, Cohn-Haft M, Figueiredo LFA, Straube FC, Cesari E (2015) Annotated checklist of the birds of Brazil by the Brazilian Ornithological Records Committee. *Revista Brasileira de Ornitológica* 23: 91–298.
- Pinto OMO (1938) Catálogo das Aves do Brasil e lista dos exemplares que as representam no Museu Paulista. Museu Paulista, São Paulo, 566 pp. <https://doi.org/10.5962/bhl.title.99663>
- Pinto OMO (1944) Catalogo das Aves do Brasil e Lista dos Exemplares Existentes na Coleção do Departamento de Zoologia, Vol. 2. Departamento de Zoologia, São Paulo, 700 pp.
- R Core Team (2016) R: A language and environment for statistical computing, R Foundation for Statistical Computing. Available online at: <https://www.R-project.org/> [Accessed: 18/03/2016]
- Ribeiro MC, Metzger JP, Martensen AC, Ponzoni FJ, Hirota MM (2009) The Brazilian Atlantic forest: how much is left, and how is the remaining forest distributed? Implications for conservation. *Biological Conservation* 142: 1141–1153. <https://doi.org/10.1016/j.biocon.2009.02.021>
- Ribon R (2010) Amostragem de aves pelo método de listas de Mackinnon. In: Von Matter S, Straube FC, Accordi IA, Piacentini VQ, Cândido Jr JF (Eds) *Ornitologia e conservação: ciência aplicada, técnicas de pesquisa e levantamento*. Technical Books, Rio de Janeiro, 33–44.
- Salis SM, Shepherd GJ, Joly CA (1995) Floristic comparison of mesophytic semideciduous forests of the interior of the state of São Paulo, Southeast Brazil. *Vegetatio* 119: 155–164. <https://doi.org/10.1007/BF00045595>
- São Paulo (2014) Decreto nº 60.133, de 7 de Fevereiro de 2014. Declara as espécies da fauna silvestre ameaçadas de extinção, as quase ameaçadas e as deficientes de dados para avaliação no Estado de S. Paulo e dá providências correlatas. <http://www.al.sp.gov.br/repositorio/legislacao/decreto/2014/decreto-60133-07.02.2014.html> [Accessed: 18/03/2016]
- Sick H (1997) *Ornitologia brasileira*. Editora Nova Fronteira, Rio de Janeiro, 912 pp.
- Silveira LF (2009) *Phylloscates eximius* (Temminck, 1822). In: Bresnan PM, Kierulff MCM, Sugieda AM (Eds) *Fauna ameaçada de extinção no estado de São Paulo: vertebrados*. Fundação Parque Zoológico de São Paulo, Secretaria do Meio Ambiente, São Paulo, 223.
- Simpson R, Cavarzere V, Simpson E (2012) List of documented bird species from the municipality of Ubatuba, State of São Paulo, Brazil. *Papéis Avulsos de Zoologia* 52: 233–254.
- Tabanez MF, Durigan G, Keurogkian A, Barbosa AF, Freitas CA, Silva CEF, Silva DA, Eaton EP, Brisolla G, Faria HH, Mattos IFA, Lobo MT, Barbosa MR, Rossi M, Souza MG, Machado RB, Pfeif-

- er RM, Ramos VS, Andrade WJ, Contiere WA (2005) Plano de Manejo da Estação Ecológica dos Caetetus. IF, Série Registros 29: 1–104.
- Tonetti VR, Pizo MA (2016) Density and microhabitat preference of the Southern Bristle-tyrant (*Phylloscartes eximius*): Conservation policy implications. *The Condor: Ornithological Applications* 118: 791–803. <https://doi.org/10.1650/CONDOR-16-89.1>
- Uezu A, Gaban-Lima R (2003) Anexo 3: espécie de aves registradas na região do Parque Estadual de Porto Ferreira (incluindo a estação experimental e estação ecológica de Jataí e o Parque Estadual de Vassununga). In: Tabanez MF, Zanchetta D (Eds) Plano de Manejo do Parque Estadual de Porto Ferreira. Secretaria do Meio Ambiente do Estado de São Paulo, Instituto Florestal, São Paulo, 16–23.
- Uezu A, Metzger JP (2016) Time-Lag in Responses of Birds to Atlantic Forest Fragmentation: Restoration Opportunity and Urgency. *PLoS ONE* 11: e0147909. <https://doi.org/10.1371/journal.pone.0147909>
- Vielliard JME, Almeida MEC, Anjos L, Silva WR (2010) Levantamento quantitativo por pontos de escuta e o Índice Pontual de Abundância (IPA). In: Von Matter S, Straube FC, Accordi IA, Piacentini VQ, Cândido Jr JF (Eds) Ornitologia e Conservação: Ciência Aplicada, Técnicas de Pesquisa e Levantamento. Technical Books, Rio de Janeiro, 47–06.
- Vilar DD (2007) Água aos Cântaros – Os reservatórios da Cantareira: um estudo de arqueologia industrial. PhD thesis, Universidade de São Paulo, São Paulo. Available online at: <http://www.teses.usp.br/teses/disponiveis/71/71131/tde-18032008-111130/pt-br.php> [Accessed: 19/03/2016]
- Ward Jr JH (1963) Hierarchical Grouping to Optimize an Objective Function. *Journal of the American Statistical Association* 58: 236–244. <https://doi.org/10.1080/01621459.1963.10500845>
- Whately M, Cunha P (2007) Cantareira 2006: um olhar sobre o maior manancial de água da Região Metropolitana de São Paulo. Instituto Socioambiental, São Paulo, 67 pp.
- Walther B (2016). Rufous-tailed Attila (*Attila phoenicurus*). In: del Hoyo J, Elliott A, Sargatal J, Christie DA, de Juana E (Eds) *Handbook of the Birds of the World Alive*. Lynx Edicions, Barcelona. <http://www.hbw.com/node/57521> [Accessed: 27/05/2016]
- WikiAves (2016) A encyclopédia das aves do Brasil. <http://www.wikiaves.com.br> [Accessed: 19/03/2016]
- Willis EO, Oniki Y (1981) Levantamento preliminar de aves em treze áreas do estado de São Paulo. *Revista Brasileira de Biologia* 41: 121–135. <https://doi.org/10.1007/BF00344966>
- Wolda H (1981) Similarity Indices, Sample Size and Diversity. *Oecologia* 50: 296–302.
- Xeno-Canto (2016) Xeno-Canto, compartilhando sons de aves do mundo todo. <http://www.xeno-canto.org> [Accessed: 19/03/2016]

APPENDIX 1

Species recorded in Serra da Cantareira. Bird families are in capital letters and in bold with the number of species in parenthesis. Nomenclature followed the Brazilian Ornithological Records Committee classification (Piacentini et al. 2015); ^{atl} = Atlantic Forest endemics (Lima 2013); ^{fo} = forest species (Parker et al. 1996); ^{sp} = species threatened in São Paulo state (São Paulo 2014); ^{BR} = species threatened in Brazil (MMA 2014); ^{GL} = globally threatened species (BirdLife 2016); * = birds lacking documented records (museum specimen, picture or song record); † = species without recent records (i.e. after Graham 1992); Locality = specific location where each species were recorded; ALSP = Alberto Löfgren State Park; CSP = Cantareira State Park, including the locality “estrada do lenhador”; Patch = species recorded in the inner slope patches (A1, A3, A4, A5 and the private reserve Sítio Capuavinha); Serra Cantareira = species recorded in Serra da Cantareira and where the specific location was not pointed; Source = record source; 1 = authors’ records during unsystematic surveys; 2 = systematic surveys (fixed-points[100 h sample effort]) performed by Vinicius R. Tonetti in CSP; 3 = systematic study (line transect [80 h] and mist-nest [2,600 nest-hours]) performed by Marco A. Rego along with other researches (names in Acknowledgements) in A1-A5; 4 = records from literature; 5 = museum specimens; 6 = species in WikiAves and/or Xeno-Canto databases.

Species	English Name	Locality	Source
Tinamidae (3)			
<i>Tinamus solitarius</i> (Vieillot, 1819)	Solitary Tinamou ^{atl, fo, sp}	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Crypturellus obsoletus</i> (Temminck, 1815)	Brown Tinamou ^{fo}	CSP, Patch	1, 2, 3, 4, 6
<i>Crypturellus tataupa</i> (Temminck, 1815)	Tataupa Tinamou ^{fo}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
Anatidae (4)			
<i>Cairina moschata</i> (Linnaeus, 1758)	Muscovy Duck	ALSP, CSP	6
<i>Dendrocygna viduata</i> (Linnaeus, 1766)	White-faced Whistling-Duck	ALSP, CSP	1, 4, 6
<i>Dendrocygna autumnalis</i> (Linnaeus, 1758)	Black-bellied Whistling-Duck	ALSP	4, 6
<i>Amazonetta brasiliensis</i> (Gmelin, 1789)	Brazilian Teal	ALSP, CSP, Patch	1, 3, 4, 6
Cracidae (1)			
<i>Penelope obscura</i> Temminck, 1815	Dusky-legged Guan ^{fo}	ALSP, CSP, Patch	1, 3, 4, 6



Species	English Name	Locality	Source
Odontophoridae (1)			
<i>Odontophorus capueira</i> (Spix, 1825)	Spot-winged Wood-Quail ^{atl, fo}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
Podicipedidae (2)			
<i>Tachybaptus dominicus</i> (Linnaeus, 1766)	Least Grebe	ALSP, CSP	1, 4, 6
<i>Podilymbus podiceps</i> (Linnaeus, 1758)	Pied-billed Grebe	ALSP, CSP	1, 4, 6
Phalacrocoracidae (1)			
<i>Nannopterum brasiliense</i> (Gmelin, 1789)	Neotropic Cormorant	ALSP, CSP	1, 4, 6
Anhingidae (1)			
<i>Anhinga anhinga</i> (Linnaeus, 1766)	Anhinga	CSP	1, 6
Ardeidae (7)			
<i>Nycticorax nycticorax</i> (Linnaeus, 1758)	Black-crowned Night-Heron	ALSP, CSP	1, 4, 6
<i>Butorides striata</i> (Linnaeus, 1758)	Striated Heron ^{fo}	ALSP, CSP	1, 4, 6
<i>Bubulcus ibis</i> (Linnaeus, 1758)	Cattle Egret	ALSP, Patch	3, 6
<i>Ardea cocoi</i> Linnaeus, 1766	Coco Heron	ALSP, CSP	1, 4, 6
<i>Ardea alba</i> Linnaeus, 1758	Great Egret	ALSP, CSP, Patch	1, 3, 4, 6
<i>Syrigma sibilatrix</i> (Temminck, 1824)	Whistling Heron*	CSP	1, 4
<i>Egretta thula</i> (Molina, 1782)	Snowy Egret	ALSP, CSP, Patch	1, 3, 4, 6
Threskiornithidae (1)			
<i>Mesembrinibis cayennensis</i> (Gmelin, 1789)	Green Ibis ^{fo}	ALSP, CSP, Patch	1, 3, 4, 6
Cathartidae (3)			
<i>Cathartes aura</i> (Linnaeus, 1758)	Turkey Vulture ^{fo}	CSP	1, 4, 6
<i>Coragyps atratus</i> (Bechstein, 1793)	Black Vulture ^{fo}	ALSP, CSP, Patch	1, 3, 4, 6
<i>Sarcoramphus papa</i> (Linnaeus, 1758)	King Vulture ^{fo, sp}	CSP	6
Accipitridae (17)			
<i>Leptodon cayanensis</i> (Latham, 1790)	Gray-headed Kite ^{fo}	ALSP, CSP, Patch	1, 3, 4, 6
<i>Chondrohierax uncinatus</i> (Temminck, 1822)	Hook-billed Kite ^{fo}	CSP	1, 5
<i>Elanoides forficatus</i> (Linnaeus, 1758)	Swallow-tailed Kite ^{fo}	CSP	1, 6
<i>Harpagus diodon</i> (Temminck, 1823)	Rufous-thighed Kite ^{fo}	ALSP, CSP	1, 4, 6
<i>Accipiter poliogaster</i> (Temminck, 1824)	Gray-bellied Hawk ^{fo}	CSP	1, 2
<i>Accipiter striatus</i> Vieillot, 1808	Sharp-shinned Hawk ^{fo}	ALSP, CSP	1, 4, 6
<i>Ictinia plumbea</i> (Gmelin, 1788)	Plumbeous Kite ^{fo}	CSP	1, 2, 4, 6
<i>Rostrhamus sociabilis</i> (Vieillot, 1817)	Snail Kite *	CSP, Patch	1, 3
<i>Geranospiza caerulescens</i> (Vieillot, 1817)	Crane Hawk ^{fo}	CSP	6
<i>Heterospizias meridionalis</i> (Latham, 1790)	Savanna Hawk *	CSP	1
<i>Amadonastur lacernulatus</i> (Temminck, 1827)	White-necked Hawk ^{atl, fo, sp, br, gl}	ALSP, CSP	1, 4, 5, 6
<i>Rupornis magnirostris</i> (Gmelin, 1788)	Roadside Hawk ^{fo}	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Parabuteo unicinctus</i> (Temminck, 1824)	Harris's Hawk ^{fo, sp * ‡}	Serra Cantareira	4
<i>Parabuteo leucorrhous</i> (Quoy & Gaimard, 1824)	White-rumped Hawk ^{fo *}	CSP	1, 4
<i>Geranoaetus albicaudatus</i> (Vieillot, 1816)	White-tailed Hawk	CSP, Patch	1, 3, 4, 6
<i>Buteo brachyurus</i> Vieillot, 1816	Short-tailed Hawk ^{fo}	ALSP, CSP, Patch	1, 3, 4, 6
<i>Spizaetus tyrannus</i> (Wied, 1820)	Black Hawk-Eagle ^{fo, sp}	CSP, Patch	1, 3, 4, 6
Rallidae (5)			
<i>Aramides cajaneus</i> (Statius Muller, 1776)	Gray-necked Wood-Rail	CSP	1, 6
<i>Aramides saracura</i> (Spix, 1825)	Slaty-breasted Wood-Rail ^{atl}	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Lateralus melanophaius</i> (Vieillot, 1819)	Rufous-sided Crake *	ALSP, CSP	1, 4
<i>Pardirallus nigricans</i> (Vieillot, 1819)	Blackish Rail	ALSP, CSP	1, 4, 6
<i>Gallinula galeata</i> (Lichtenstein, 1818)	Common Gallinule	ALSP, CSP	1, 4, 6
Charadriidae (1)			
<i>Vanellus chilensis</i> (Molina, 1782)	Southern Lapwing	ALSP, CSP, Patch	1, 3, 4, 6
Scolopacidae (1)			
<i>Tringa solitaria</i> Wilson, 1813	Solitary Sandpiper	CSP	1, 6



Species	English Name	Locality	Source
Jacanidae (1)			
<i>Jacana jacana</i> (Linnaeus, 1766)	Wattled Jacana	ALSP, Patch	1, 3, 4, 6
Columbidae (12)			
<i>Columbina talpacoti</i> (Temminck, 1811)	Ruddy Ground-Dove	ALSP, CSP, Patch	1, 3, 4, 6
<i>Columbina squammata</i> (Lesson, 1831)	Scaled Dove *	ALSP	4
<i>Claravis pretiosa</i> (Ferrari-Perez, 1886)	Blue Ground-Dove ^{FO}	CSP	1, 6
<i>Claravis geoffroyi</i> (Temminck, 1811)	Purple-winged Ground-Dove ^{ATL, FO} SP, BR, GL, ♀	CSP	5
<i>Columba livia</i> Gmelin, 1789	Rock Pigeon ^{Exo}	ALSP, CSP, Patch	1, 4, 6
<i>Patagioenas picazuro</i> (Temminck, 1813)	Picazuro Pigeon ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Patagioenas cayennensis</i> (Bonnaterre, 1792)	Pale-vented Pigeon ^{FO *}	CSP, Patch	1, 2, 4
<i>Patagioenas plumbea</i> (Vieillot, 1818)	Plumbeous Pigeon ^{FO}	CSP, Patch	1, 2, 3, 4, 6
<i>Zenaida auriculata</i> (Des Murs, 1847)	Eared Dove *	CSP, Patch	1, 3, 4
<i>Leptotila verreauxi</i> Bonaparte, 1855	White-tipped Dove ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Leptotila rufaxilla</i> (Richard & Bernard, 1792)	Gray-fronted Dove ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Geotrygon montana</i> (Linnaeus, 1758)	Ruddy Quail-Dove ^{FO}	CSP, Patch	1, 2, 3, 4, 6
Cuculidae (8)			
<i>Piaya cayana</i> (Linnaeus, 1766)	Squirrel Cuckoo ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
<i>Coccyzus melacoryphus</i> Vieillot, 1817	Dark-billed Cuckoo ^{FO *}	CSP	4
<i>Coccyzus americanus</i> (Linnaeus, 1758)	Yellow-billed Cuckoo ^{FO}	CSP	1, 4, 6
<i>Coccyzus euleri</i> Cabanis, 1873	Pearly-breasted Cuckoo ^{FO}	CSP	4, 6
<i>Crotophaga ani</i> Linnaeus, 1758	Smooth-billed Ani	ALSP, CSP, Patch	1, 3, 4, 5, 6
<i>Guira guira</i> (Gmelin, 1788)	Guira Cuckoo *	ALSP, CSP, Patch	1, 4
<i>Tapera naevia</i> (Linnaeus, 1766)	Striped Cuckoo *	CSP, Patch	3
<i>Dromococcyx pavoninus</i> Pelzeln, 1870	Pavonine Cuckoo ^{FO *}	CSP, Patch	1, 2, 3
Tytonidae (1)			
<i>Tyto furcata</i> (Temminck, 1827)	American Barn Owl *	ALSP, CSP	1, 4
Strigidae (6)			
<i>Megascops choliba</i> (Vieillot, 1817)	Tropical Screech-Owl ^{FO}	ALSP, CSP, Patch	1, 3, 4, 6
<i>Pulsatrix koeniswaldiana</i> (Bertoni & Bertoni, 1901)	Tawny-browed Owl ^{ATL, FO}	ALSP, CSP, Patch	1, 3, 4, 5, 6
<i>Strix hylophila</i> Temminck, 1825	Rusty-barred Owl ^{ATL, FO *}	CSP	1, 3
<i>Strix virgata</i> (Cassin, 1849)	Mottled Owl ^{FO}	ALSP, CSP, Patch	1, 3, 4, 5
<i>Athene cunicularia</i> (Molina, 1782)	Burrowing Owl *	ALSP, Patch	1, 3, 4
<i>Asio clamator</i> (Vieillot, 1808)	Striped Owl	ALSP, CSP	4, 5, 6
Nyctibiidae (1)			
<i>Nyctibius griseus</i> (Gmelin, 1789)	Common Potoo ^{FO *}	CSP	1, 4
Caprimulgidae (7)			
<i>Nyctiphrynus ocellatus</i> (Tschudi, 1844)	Ocellated Poorwill ^{FO}	CSP, Patch	1, 3
<i>Lurocalis semitorquatus</i> (Gmelin, 1789)	Short-tailed Nighthawk ^{FO}	ALSP, CSP, Patch	1, 3, 4, 6
<i>Nyctidromus albicollis</i> (Gmelin, 1789)	Pauraque ^{FO}	ALSP, CSP, Patch	1, 3, 4, 6
<i>Hydropsalis parvula</i> (Gould, 1837)	Little Nightjar ^{FO *}	Patch	1, 3
<i>Hydropsalis longirostris</i> (Bonaparte, 1825)	Band-winged Nightjar	CSP	6
<i>Hydropsalis torquata</i> (Gmelin, 1789)	Scissor-tailed Nightjar ^{FO *}	CSP, Patch	1, 3, 4
<i>Hydropsalis forcipata</i> (Nitzsch, 1840)	Long-trained Nightjar ^{ATL, FO *}	CSP, Patch	3, 4
Apodidae (2)			
<i>Streptoprocne zonaris</i> (Shaw, 1796)	White-collared Swift ^{FO *}	ALSP, CSP, Patch	1, 4
<i>Chaetura meridionalis</i> Hellmayr, 1907	Sick's Swift ^{FO}	ALSP, CSP, Patch	1, 3, 4, 6
Trochilidae (14)			
<i>Phaethornis pretrei</i> (Lesson & Delattre, 1839)	Planalto Hermit ^{FO *}	ALSP, CSP, Patch	1, 3, 4
<i>Phaethornis eurynome</i> (Lesson, 1832)	Scale-throated Hermit ^{ATL, FO}	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Eupetomena macroura</i> (Gmelin, 1788)	Swallow-tailed Hummingbird	ALSP, CSP, Patch	1, 4, 6



Species	English Name	Locality	Source
<i>Aphantochroa cirrochloris</i> (Vieillot, 1818)	Sombre Hummingbird ^{FO}	ALSP, CSP, Patch	1, 4, 5, 6
<i>Florisuga fusca</i> (Vieillot, 1817)	Black Jacobin ^{FO}	ALSP, CSP, Patch	1, 2, 4, 5, 6
<i>Anthracothorax nigricollis</i> (Vieillot, 1817)	Black-throated Mango ^{FO}	CSP	6
<i>Lophornis chalybeus</i> (Vieillot, 1822)	Festive Coquette ^{FO}	ALSP, CSP	1, 4, 6
<i>Chlorostilbon lucidus</i> (Shaw, 1812)	Glittering-bellied Emerald ^{FO}	ALSP, CSP, Patch	1, 3, 4, 5, 6
<i>Thalurania glaukopis</i> (Gmelin, 1788)	Violet-capped Woodnymph ^{ATL, FO}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
<i>Leucochloris albicollis</i> (Vieillot, 1818)	White-throated Hummingbird ^{FO}	ALSP, CSP, Patch	1, 4, 6
<i>Amazilia versicolor</i> (Vieillot, 1818)	Versicolored Emerald ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Amazilia fimbriata</i> (Gmelin, 1788)	Glittering-throated Emerald ^{FO}	CSP	1, 2, 4, 6
<i>Amazilia lactea</i> (Lesson, 1832)	Sapphire-spangled Emerald ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Heliodoxa rubricauda</i> (Boddaert, 1783)	Brazilian Ruby ^{ATL, FO}	ALSP, CSP	1, 4, 5, 6
Trogonidae (2)			
<i>Trogon viridis</i> Linnaeus, 1766	White-tailed Tropicbird ^{FO}	ALSP, CSP	5, 6
<i>Trogon surrucura</i> Vieillot, 1817	Surucua Tropicbird ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
Alcedinidae (3)			
<i>Megaceryle torquata</i> (Linnaeus, 1766)	Ringed Kingfisher	ALSP, CSP, Patch	1, 3, 4, 6
<i>Chloroceryle amazona</i> (Latham, 1790)	Amazon Kingfisher	ALSP, CSP	1, 3, 4, 6
<i>Chloroceryle americana</i> (Gmelin, 1788)	Green Kingfisher	ALSP, CSP, Patch	1, 3, 4, 6
Momotidae (1)			
<i>Baryphthengus ruficapillus</i> (Vieillot, 1818)	Rufous-capped Motmot ^{FO *}	Patch	1, 3
Bucconidae (1)			
<i>Malacoptila striata</i> (Spix, 1824)	Crescent-chested Puffbird ^{ATL, FO}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
Ramphastidae (5)			
<i>Ramphastos toco</i> Statius Muller, 1776	Toco Toucan	ALSP, CSP	6
<i>Ramphastos vitellinus</i> Lichtenstein, 1823	Channel-billed Toucan ^{FO, GL}	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Ramphastos dicolorus</i> Linnaeus, 1766	Red-breasted Toucan ^{ATL, FO}	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Selenidera maculirostris</i> (Lichtenstein, 1823)	Spot-billed Toucanet ^{ATL, FO, SP}	Patch	1, 3, 5
<i>Pteroglossus bailloni</i> (Vieillot, 1819)	Saffron Toucanet ^{ATL, FO, SP}	ALSP, CSP	1, 4, 5, 6
Picidae (11)			
<i>Picumnus cirratus</i> Temminck, 1825	White-barred Piculet ^{FO *}	Patch	1
<i>Picumnus temminckii</i> Lafresnaye, 1845	Ochre-collared Piculet ^{ATL, FO}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
<i>Melanerpes candidus</i> (Otto, 1796)	White Woodpecker *	ALSP, CSP, Patch	1, 3, 4
<i>Melanerpes flavifrons</i> (Vieillot, 1818)	Yellow-fronted Woodpecker ^{FO}	CSP	1, 3, 4, 5
<i>Veniliornis spilogaster</i> (Wagler, 1827)	White-spotted Woodpecker ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Piculus aurulentus</i> (Temminck, 1821)	Yellow-browed Woodpecker ^{ATL, FO}	CSP, Patch	1, 3, 4, 6
<i>Colaptes melanochloros</i> (Gmelin, 1788)	Green-barred Woodpecker ^{FO}	ALSP, CSP, Patch	1, 3, 4, 6
<i>Colaptes campestris</i> (Vieillot, 1818)	Campo Flicker *	ALSP, CSP, Patch	1, 3, 4
<i>Celeus flavescens</i> (Gmelin, 1788)	Blond-crested Woodpecker ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Dryocopus lineatus</i> (Linnaeus, 1766)	Lineated Woodpecker ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Campephilus robustus</i> (Lichtenstein, 1818)	Robust Woodpecker ^{ATL, FO *}	CSP, Patch	1, 2, 3, 4
Cariamidae (1)			
<i>Cariama cristata</i> (Linnaeus, 1766)	Red-legged Seriema *	CSP, Patch	1, 3, 4
Falconidae (6)			
<i>Caracara plancus</i> (Miller, 1777)	Southern Caracara	ALSP, CSP, Patch	1, 3, 4, 6
<i>Milvago chimachima</i> (Vieillot, 1816)	Yellow-headed Caracara	ALSP, CSP, Patch	1, 3, 4, 6
<i>Herpetotheres cachinnans</i> (Linnaeus, 1758)	Laughing Falcon ^{FO}	CSP, Patch	1, 2, 3, 4, 6
<i>Micrastur ruficollis</i> (Vieillot, 1817)	Barred Forest-Falcon ^{FO}	CAP, Patch	1, 2, 3, 4, 6
<i>Micrastur semitorquatus</i> (Vieillot, 1817)	Collared Forest-Falcon ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Falco femoralis</i> Temminck, 1822	Aplomado Falcon	CSP	1, 6
Psittacidae (8)			
<i>Diopsittaca nobilis</i> (Linnaeus, 1758)	Red-shouldered Macaw ^{EXO, FO, SP}	ALSP, CSP	1, 4, 6



Species	English Name	Locality	Source
<i>Psittacula leucophthalmus</i> (Statius Muller, 1776)	White-eyed Parakeet ^{FO *}	CSP, Patch	1, 2, 3, 4
<i>Pyrrhura frontalis</i> (Vieillot, 1817)	Maroon-bellied Parakeet ^{atl, FO}	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Forpus xanthopterygius</i> (Spix, 1824)	Blue-winged Parrotlet ^{FO}	ALSP, CSP, Patch	1, 3, 4, 6
<i>Brotogeris tirica</i> (Gmelin, 1788)	Plain Parakeet ^{atl, FO}	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Pionopsitta pileata</i> (Scopoli, 1769)	Red-capped Parrot ^{atl, FO}	ALSP, CSP	1, 4
<i>Pionus maximiliani</i> (Kuhl, 1820)	Scaly-headed Parrot ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Amazona aestiva</i> (Linnaeus, 1758)	Blue-fronted Parrot ^{Exo}	ALSP, CSP	1, 2, 4, 6
Thamnophilidae (15)			
<i>Rhopias gularis</i> (Spix, 1825)	Star-throated Antwren ^{atl}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
<i>Dysithamnus stictothorax</i> (Temminck, 1823)	Spot-breasted Antvireo ^{atl, FO ‡}	—	5
<i>Dysithamnus mentalis</i> (Temminck, 1823)	Plain Antvireo ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
<i>Herpsilochmus rufimarginatus</i> (Temminck, 1822)	Rufous-winged Antwren ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
<i>Thamnophilus doliatus</i> (Linnaeus, 1764)	Barred Antshrike	ALSP	4, 6
<i>Thamnophilus ruficapillus</i> Vieillot, 1816	Rufous-capped Antshrike	ALSP, CSP, Patch	4, 6
<i>Thamnophilus caerulescens</i> Vieillot, 1816	Variable Antshrike ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
<i>Hypoedaleus guttatus</i> (Vieillot, 1816)	Spot-backed Antshrike ^{atl, FO}	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Batara cinerea</i> (Vieillot, 1819)	Giant Antshrike ^{FO}	CSP, Patch	1, 2, 3, 4, 6
<i>Mackenziaena leachii</i> (Such, 1825)	Large-tailed Antshrike ^{atl, FO *}	CSP, Patch	1, 3, 4
<i>Mackenziaena severa</i> (Lichtenstein, 1823)	Tufted Antshrike ^{atl, FO *}	Patch	3
<i>Myrmotherula squamosus</i> (Pelzeln, 1868)	Squamate Antbird ^{atl, FO}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
<i>Pyriglenia leucoptera</i> (Vieillot, 1818)	White-shouldered Fire-eye ^{atl, FO}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
<i>Drymophila ferruginea</i> (Temminck, 1822)	Ferruginous Antbird ^{atl, FO}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
<i>Drymophila ochropyga</i> (Hellmayr, 1906)	Ochre-rumped Antbird ^{atl, FO, SP}	CSP, Patch	1, 2, 3, 4, 5, 6
Conopophagidae (1)			
<i>Conopophaga lineata</i> (Wied, 1831)	Rufous Gnat-eater ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
Grallariidae (2)			
<i>Grallaria varia</i> (Boddaert, 1783)	Variegated Antpitta ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
<i>Hylopezus nattereri</i> (Pinto, 1937)	Speckle-breasted Antpitta ^{atl, FO ‡}	—	5
Rhinocryptidae (2)			
<i>Eleoscytalopus indigoticus</i> (Wied, 1831)	White-breasted Tapaculo ^{atl, FO *}	CSP	1
<i>Scytalopus speluncae</i> (Ménétriès, 1835)	Mouse-colored Tapaculo ^{atl, FO *}	CSP	1, 4
Formicariidae (2)			
<i>Chamaezza campanisona</i> (Lichtenstein, 1823)	Short-tailed Antthrush ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 5
<i>Chamaezza meruloides</i> Vigors, 1825	Such's Antthrush ^{atl, FO}	CSP, Patch	1, 2, 3, 4, 5, 6
Scleruridae (1)			
<i>Sclerurus scansor</i> (Ménétriès, 1835)	Rufous-breasted Leaf-tosser ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
Dendrocolaptidae (6)			
<i>Sittasomus griseicapillus</i> (Vieillot, 1818)	Olivaceous Woodcreeper ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
<i>Xiphorhynchus fuscus</i> (Vieillot, 1818)	Lesser Woodcreeper ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
<i>Lepidocolaptes falcinellus</i> (Cabanis & Heine, 1859)	Scalloped Woodcreeper ^{atl, FO}	CSP, Patch	1, 3, 4, 5, 6
<i>Dendrocolaptes platyrostris</i> Spix, 1825	Planalto Woodcreeper ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
<i>Xiphocolaptes albicollis</i> (Vieillot, 1818)	White-throated Woodcreeper ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 6
Xenopidae (2)			
<i>Xenops minutus</i> (Sparrman, 1788)	Plain Xenops ^{FO}	CSP, Patch	1, 2, 3, 4, 6
<i>Xenops rutilans</i> Temminck, 1821	Streaked Xenops ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
Furnariidae (18)			
<i>Furnarius figulus</i> (Lichtenstein, 1823)	Wing-banded Hornero	ALSP, CSP	1, 6
<i>Furnarius rufus</i> (Gmelin, 1788)	Rufous Hornero	ALSP, CSP, Patch	1, 3, 4, 6
<i>Lochmias nematura</i> (Lichtenstein, 1823)	Sharp-tailed Streamcreeper ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
<i>Automolus leucophthalmus</i> (Wied, 1821)	White-eyed Foliage-gleaner ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
<i>Anabazenops fuscus</i> (Vieillot, 1816)	White-collared Foliage-gleaner ^{atl, FO}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6



Species	English Name	Locality	Source
<i>Anabacerthia amaroensis</i> (Temminck, 1823)	White-browed Foliage-gleaner ^{atl, fo}	CSP, Patch	1, 2, 3, 4, 6
<i>Philydor atricapillus</i> (Wied, 1821)	Black-capped Foliage-gleaner ^{atl, fo}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
<i>Philydor rufum</i> (Vieillot, 1818)	Buff-fronted Foliage-gleaner ^{fo}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
<i>Heliobletus contaminatus</i> Berlepsch, 1885	Sharp-billed Treecrusher ^{atl, fo}	CSP, Patch	1, 2, 3, 4, 6
<i>Syndactyla rufosuperciliata</i> (Lafresnaye, 1832)	Buff-browed Foliage-gleaner ^{fo}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
<i>Phacellodomus erythrophthalmus</i> (Wied, 1821)	Orange-eyed Thornbird ^{atl, fo}	CSP, Patch	1, 2, 3, 6
<i>Phacellodomus ferrugineigula</i> (Pelzeln, 1858)	Orange-breasted Thornbird ^{atl}	CSP, Patch	1, 3, 4, 6
<i>Certhiaxis cinnamomeus</i> (Gmelin, 1788)	Yellow-chinned Spinetail	ALSP, Patch	1, 3, 4, 6
<i>Synallaxis ruficapilla</i> Vieillot, 1819	Rufous-capped Spinetail ^{atl, fo}	CSP, Patch	1, 2, 3, 4, 5, 6
<i>Synallaxis cinerascens</i> Temminck, 1823	Gray-bellied Spinetail ^{fo}	CAP, Patch	1, 2, 3, 4, 5, 6
<i>Synallaxis spixi</i> Sclater, 1856	Spix's Spinetail	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Cranioleuca pallida</i> (Wied, 1831)	Pallid Spinetail ^{atl, fo}	ALSP, CSP, Patch	1, 2, 3, 4, 6
Pipridae (4)			
<i>Neopelma chrysophorum</i> Pinto, 1944	Serra do Mar Tyrant-Manakin ^{atl, fo, ♀}	Serra Cantareira	5
<i>Manacus manacus</i> (Linnaeus, 1766)	White-bearded Manakin ^{fo, ♀ *}	Serra Cantareira	5
<i>Ilicura militaris</i> (Shaw & Nodder, 1809)	Pin-tailed Manakin ^{atl, fo}	CSP	1, 2, 4, 6
<i>Chiroxiphia caudata</i> (Shaw & Nodder, 1793)	Swallow-tailed Manakin ^{atl, fo}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
Oxyruncidae (1)			
<i>Oxyruncus cristatus</i> Swainson, 1821	Sharpbill	CSP	1, 4, 6
ONYCHORHYNCHIDAE (1)			
<i>Myioobius atricaudus</i> Lawrence, 1863	Black-tailed Flycatcher ^{fo, ♀}	Serra Cantareira	5
Tityridae (6)			
<i>Schiffornis virescens</i> (Lafresnaye, 1838)	Greenish Schiffornis ^{fo}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
<i>Laniisoma elegans</i> (Thunberg, 1823)	Shrike-like Cotinga ^{atl, fo, sp}	CSP	6
<i>Tityra cayana</i> (Linnaeus, 1766)	Black-tailed Tityra	ALSP, CSP	1, 4, 6
<i>Pachyramphus castaneus</i> (Jardine & Selby, 1827)	Chestnut-crowned Becard ^{fo}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
<i>Pachyramphus polychropterus</i> (Vieillot, 1818)	White-winged Becard ^{fo}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
<i>Pachyramphus validus</i> (Lichtenstein, 1823)	Crested Becard ^{fo}	ALSP, CSP	1, 2, 4, 6
Cotingidae (4)			
<i>Phibalura flavirostris</i> Vieillot, 1816	Swallow-tailed Cotinga ^{atl, fo}	CSP	1, 4
<i>Pyroderus scutatus</i> (Shaw, 1792)	Red-ruffed Fruitcrow ^{fo, sp}	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Lipaugus lanioides</i> (Lesson, 1844)	Cinnamon-vented Piha ^{atl *}	CSP	1
<i>Procnias nudicollis</i> (Vieillot, 1817)	Bare-throated Bellbird ^{atl, fo, sp, gl}	ALSP, CSP	1, 4, 6
Platyrinchidae (1)			
<i>Platyrinchus mystaceus</i> Vieillot, 1818	White-throated Spadebill ^{fo}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
Rhynchocyclidae (13)			
<i>Mionectes rufiventris</i> Cabanis, 1846	Gray-hooded Flycatcher ^{fo}	CSP, Patch	1, 2, 3, 4, 5, 6
<i>Leptopogon amaurocephalus</i> Tschudi, 1846	Sepia-capped Flycatcher ^{fo}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
<i>Corythopis delalandi</i> (Lesson, 1830)	Southern Antpipit ^{fo}	CSP	3
<i>Phylloscartes eximius</i> (Temminck, 1822)	Southern Bristle-Tyrant ^{atl, fo, sp}	CSP, Patch	1, 2, 3, 4, 6
<i>Phylloscartes ventralis</i> (Temminck, 1824)	Mottle-cheeked Tyrannulet ^{fo}	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Tolmomyias sulphurescens</i> (Spix, 1825)	Yellow-olive Flycatcher ^{fo}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
<i>Todirostrum poliocephalum</i> (Wied, 1831)	Yellow-lored Tody-Flycatcher ^{atl, fo}	ALSP, CSP	1, 2, 4, 5, 6
<i>Todirostrum cinereum</i> (Linnaeus, 1766)	Common Tody-Flycatcher	ALSP, CSP	1, 2, 4, 6
<i>Poecilotriccus plumbeiceps</i> (Lafresnaye, 1846)	Ochre-faced Tody-Flycatcher ^{fo}	CSP, Patch	1, 2, 3, 4, 6
<i>Myiornis auricularis</i> (Vieillot, 1818)	Eared Pygmy-Tyrant ^{atl, fo}	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Hemitriccus diops</i> (Temminck, 1822)	Drab-breasted Pygmy-Tyrant ^{atl, fo}	CSP, Patch	1, 2, 3, 4, 5, 6
<i>Hemitriccus orbitatus</i> (Wied, 1831)	Eye-ringed Tody-Tyrant ^{atl, fo}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
<i>Hemitriccus nidipendulus</i> (Wied, 1831)	Hangnest Tody-Tyrant ^{atl, fo}	CSP, Patch	1, 2, 3, 4, 5
Tyrannidae (35)			
<i>Hirundinea ferruginea</i> (Gmelin, 1788)	Cliff Flycatcher *	CSP, Patch	4



Species	English Name	Locality	Source
<i>Tyranniscus burmeisteri</i> (Cabanis & Heine, 1859)	Rough-legged Tyrannulet ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Camptostoma obsoletum</i> (Temminck, 1824)	Southern Beardless-Tyrannulet	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Elaenia flavogaster</i> (Thunberg, 1822)	Yellow-bellied Elaenia	ALSP, CSP, Patch	1, 4, 6
<i>Elaenia parvirostris</i> Pelzeln, 1868	Small-billed Elaenia ^{FO *}	CSP	1, 4
<i>Elaenia mesoleuca</i> (Deppe, 1830)	Olivaceous Elaenia ^{FO}	CSP, Patch	1, 3, 4, 5, 6
<i>Elaenia obscura</i> (d'Orbigny & Lafresnaye, 1837)	Highland Elaenia ^{FO}	Serra Cantareira	5
<i>Myiopagis caniceps</i> (Swainson, 1835)	Gray Elaenia ^{FO *}	Patch	3
<i>Phyllosmyias virescens</i> (Temminck, 1824)	Greenish Tyrannulet ^{ATL, FO}	CSP	1, 6
<i>Phyllosmyias fasciatus</i> (Thunberg, 1822)	Planalto Tyrannulet ^{FO}	ALSP, CSP, Patch	1, 3, 4
<i>Phyllosmyias griseocapilla</i> Sclater, 1862	Gray-capped Tyrannulet ^{ATL, FO}	CSP, Patch	1, 2, 3, 4
<i>Serophaea subcristata</i> (Vieillot, 1817)	White-crested Tyrannulet ^{FO *}	ALSP, CSP, Patch	1, 3, 4
<i>Attila phoenicurus</i> Pelzeln, 1868	Rufous-tailed Attila ^{FO}	CSP	4
<i>Attila rufus</i> (Vieillot, 1819)	Gray-hooded Attila ^{ATL, FO}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
<i>Legatus leucophaius</i> (Vieillot, 1818)	Piratic Flycatcher ^{FO}	ALSP, CSP, Patch	1, 2, 4, 6
<i>Ramphotrigon megacephalum</i> (Swainson, 1835)	Large-headed Flatbill ^{FO}	CSP, Patch	3, 6
<i>Myiarchus swainsoni</i> Cabanis & Heine, 1859	Swainson's Flycatcher ^{FO}	CSP, Patch	1, 2, 3, 4, 6
<i>Myiarchus ferox</i> (Gmelin, 1789)	Short-crested Flycatcher ^{FO}	ALSP, CSP, Patch	1, 3, 4, 5, 6
<i>Myiarchus tyrannulus</i> (Statius Muller, 1776)	Brown-crested Flycatcher [*]	Serra Cantareira	4
<i>Pitangus sulphuratus</i> (Linnaeus, 1766)	Great Kiskadee	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Machetornis rixosa</i> (Vieillot, 1819)	Cattle Tyrant [*]	ALSP, CSP, Patch	1, 3, 4
<i>Myiodynastes maculatus</i> (Statius Muller, 1776)	Streaked Flycatcher ^{FO}	ALSP, CSP	1, 2, 4, 6
<i>Megarynchus pitangua</i> (Linnaeus, 1766)	Boat-billed Flycatcher ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Myiozetetes similis</i> (Spix, 1825)	Social Flycatcher ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Tyrannus melancholicus</i> Vieillot, 1819	Tropical Kingbird	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Tyrannus savana</i> Vieillot, 1808	Fork-tailed Flycatcher [*]	ALSP, CSP, Patch	4
<i>Empidonax varius</i> (Vieillot, 1818)	Variegated Flycatcher ^{FO}	ALSP, CSP, Patch	1, 4, 6
<i>Colonia colonus</i> (Vieillot, 1818)	Long-tailed Tyrant ^{FO}	CSP	1, 4, 6
<i>Myiophobus fasciatus</i> (Statius Muller, 1776)	Bran-colored Flycatcher	CSP, Patch	1, 3, 4, 6
<i>Fluvicola nengeta</i> (Linnaeus, 1766)	Masked Water-Tyrant	ALSP, CSP, Patch	1, 3, 4, 6
<i>Cnemotriccus fuscatus</i> (Wied, 1831)	Fuscous Flycatcher ^{FO}	CSP, Patch	3, 4, 6
<i>Lathrotriccus euleri</i> (Cabanis, 1868)	Euler's Flycatcher ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Contopus cinereus</i> (Spix, 1825)	Tropical Pewee ^{FO}	CSP	1, 2, 4, 6
<i>Knipolegus cyanirostris</i> (Vieillot, 1818)	Blue-billed Black-Tyrant [*]	CSP	4
<i>Muscicapa vetula</i> (Lichtenstein, 1823)	Shear-tailed Gray Tyrant ^{ATL, FO}	CSP, Patch	1, 3
Vireonidae (3)			
<i>Cyclarhis gujanensis</i> (Gmelin, 1789)	Rufous-browed Peppershrike ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
<i>Hylophilus poicilotis</i> Temminck, 1822	Rufous-crowned Greenlet ^{ATL, FO}	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Vireo chivi</i> (Vieillot, 1817)	Chivi Vireo ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
Corvidae (1)			
<i>Cyanocorax cristatellus</i> (Temminck, 1823)	Curl-crested Jay	Patch	1, 3, 6
Hirundinidae (4)			
<i>Pygochelidon cyanoleuca</i> (Vieillot, 1817)	Blue-and-white Swallow	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Stelgidopteryx ruficollis</i> (Vieillot, 1817)	Southern Rough-winged Swallow	ALSP, CSP, Patch	1, 3, 4, 6
<i>Progne tapera</i> (Vieillot, 1817)	Brown-chested Martin [*]	CSP	1, 3
<i>Tachycineta leucorrhoa</i> (Vieillot, 1817)	White-rumped Swallow [*]	CSP, Patch	1, 3
Troglodytidae (1)			
<i>Troglodytes musculus</i> Naumann, 1823	Southern House Wren	ALSP, CSP, Patch	1, 2, 3, 4, 6
Donacobiidae (1)			
<i>Donacobius atricapilla</i> (Linnaeus, 1766)	Black-capped Donacobius [*]	ALSP	4
Polioptilidae (1)			
<i>Polioptila dumicola</i> (Vieillot, 1817)	Masked Gnatcatcher [*]	ALSP	4



Species	English Name	Locality	Source
Turdidae (6)			
<i>Turdus flavipes</i> Vieillot, 1818	Yellow-legged Thrush ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
<i>Turdus leucomelas</i> Vieillot, 1818	Pale-breasted Thrush ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Turdus rufiventris</i> Vieillot, 1818	Rufous-bellied Thrush ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
<i>Turdus amaurochalinus</i> Cabanis, 1850	Creamy-bellied Thrush ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
<i>Turdus subalaris</i> (Seebold, 1887)	Eastern Slaty Thrush ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Turdus albicollis</i> Vieillot, 1818	White-necked Thrush ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
Mimidae (1)			
<i>Mimus saturninus</i> (Lichtenstein, 1823)	Chalk-browed Mockingbird	ALSP, CSP, Patch	1, 3, 4, 6
Passerellidae (2)			
<i>Zonotrichia capensis</i> (Statius Muller, 1776)	Rufous-collared Sparrow	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
<i>Arremon semitorquatus</i> Swainson, 1838	Half-collared Sparrow ^{atl, FO}	CSP	1, 2, 4, 6
Parulidae (5)			
<i>Setophaga pitayumi</i> (Vieillot, 1817)	Tropical Parula ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Geothlypis aequinoctialis</i> (Gmelin, 1789)	Masked Yellowthroat	ALSP, CSP, Patch	1, 3, 4, 6
<i>Basileuterus culicivorus</i> (Deppe, 1830)	Golden-crowned Warbler ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
<i>Myiothlypis leucoblephara</i> (Vieillot, 1817)	White-browed Warbler ^{atl, FO}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
<i>Myiothlypis rivularis</i> (Wied, 1821)	Neotropical River Warbler ^{atl, FO}	CSP	6
Icteridae (6)			
<i>Psarocolius decumanus</i> (Pallas, 1769)	Crested Oropendola ^{FO *}	CSP	1, 4
<i>Cacicus chrysopterus</i> (Vigors, 1825)	Golden-winged Cacique ^{FO}	ALSP, CSP	1, 2, 4, 6
<i>Cacicus haemorrhous</i> (Linnaeus, 1766)	Red-rumped Cacique ^{FO}	ALSP, CSP	1, 4, 6
<i>Cacicus cela</i> (Linnaeus, 1758)	Yellow-rumped Cacique ^{Exo, FO}	CSP	1, 6
<i>Chrysomus ruficapillus</i> (Vieillot, 1819)	Chestnut-capped Blackbird	ALSP, Patch	4, 6
<i>Molothrus bonariensis</i> (Gmelin, 1789)	Shiny Cowbird	ALSP, CSP	1, 4, 6
Mitrospingidae (1)			
<i>Orthogonyx chloricterus</i> (Vieillot, 1819)	Olive-green Tanager ^{atl, FO}	CSP	1, 4, 6
Thraupidae (33)			
<i>Orchesticus abeillei</i> (Lesson, 1839)	Brown Tanager ^{atl, FO}	CSP	1, 2, 4, 6
<i>Pipreidea melanonota</i> (Vieillot, 1819)	Fawn-breasted Tanager ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Schistochlamys ruficapillus</i> (Vieillot, 1817)	Cinnamon Tanager ^{FO} [‡]	ALSP	4, 5
<i>Paroaria dominicana</i> (Linnaeus, 1758)	Red-cowled Cardinal ^{Exo}	CSP	6
<i>Thlypopsis sordida</i> (d'Orbigny & Lafresnaye, 1837)	Orange-headed Tanager	ALSP, CSP, Patch	1, 3, 4, 6
<i>Tangara seledon</i> (Statius Muller, 1776)	Green-headed Tanager ^{atl, FO}	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Tangara cyaniventris</i> (Vieillot, 1819)	Gilt-edged Tanager ^{atl, FO}	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Tangara desmaresti</i> (Vieillot, 1819)	Brassy-breasted Tanager ^{atl, FO}	ALSP, CSP, Patch	1, 2, 4, 5, 6
<i>Tangara sayaca</i> (Linnaeus, 1766)	Sayaca Tanager	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Tangara palmarum</i> (Wied, 1823)	Palm Tanager	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Tangara ornata</i> (Sparrman, 1789)	Golden-chevroned Tanager ^{atl, FO}	ALSP	6
<i>Tangara peruviana</i> (Desmarest, 1806)	Black-backed Tanager ^{atl, FO, SP, BR, GL}	CSP, Patch	4, 6
<i>Tangara preciosa</i> (Cabanis, 1850)	Chestnut-backed Tanager ^{FO *}	CSP, Patch	3, 4
<i>Tangara cayana</i> (Linnaeus, 1766)	Burnished-buff Tanager	ALSP, CSP, Patch	1, 3, 4, 5, 6
<i>Nemosia pileata</i> (Boddaert, 1783)	Hooded Tanager ^{FO}	ALSP, CSP	1, 6
<i>Conirostrum speciosum</i> (Temminck, 1824)	Chestnut-vented Conebill ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Sicalis flaveola</i> (Linnaeus, 1766)	Saffron Finch	CSP	4, 6
<i>Haplospiza unicolor</i> Cabanis, 1851	Uniform Finch ^{atl, FO}	CSP, Patch	1, 3, 4, 5, 6
<i>Hemithraupis ruficapilla</i> (Vieillot, 1818)	Rufous-headed Tanager ^{atl, FO}	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Volatinia jacarina</i> (Linnaeus, 1766)	Blue-black Grassquit	ALSP, CSP, Patch	1, 3, 4, 6
<i>Trichothraupis melanops</i> (Vieillot, 1818)	Black-goggled Tanager ^{FO}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
<i>Tachyphonus coronatus</i> (Vieillot, 1822)	Ruby-crowned Tanager ^{atl, FO}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
<i>Ramphocelus bresilius</i> (Linnaeus, 1766)	Brazilian Tanager ^{atl, FO}	ALSP, CSP	6



Species	English Name	Locality	Source
<i>Tersina viridis</i> (Illiger, 1811)	Swallow Tanager ^{Fo}	ALSP, CSP, Patch	1, 3, 4, 6
<i>Dacnis cayana</i> (Linnaeus, 1766)	Blue Dacnis ^{Fo}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
<i>Coereba flaveola</i> (Linnaeus, 1758)	Bananaquit	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
<i>Sporophila lineola</i> (Linnaeus, 1758)	Lined Seedeater *	ALSP, Patch	4
<i>Sporophila frontalis</i> (Verreaux, 1869)	Buffy-fronted Seedeater ^{atl, Fo, SP, BR, GL}	CSP, Patch	1, 2, 3, 4, 6
<i>Sporophila falcirostris</i> (Temminck, 1820)	Temminck's Seedeater ^{atl, Fo, SP, BR, GL}	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Sporophila caerulescens</i> (Vieillot, 1823)	Double-collared Seedeater *	ALSP, CSP, Patch	1, 3, 4
<i>Sporophila leucoptera</i> (Vieillot, 1817)	White-bellied Seedeater *	Patch	4
<i>Saltator similis</i> d'Orbigny & Lafresnaye, 1837	Green-winged Saltator ^{Fo}	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Saltator fuliginosus</i> (Daudin, 1800)	Black-throated Grosbeak ^{atl, Fo}	CSP	1, 4, 6
Cardinalidae (1)			
<i>Habia rubica</i> (Vieillot, 1817)	Red-crowned Ant-Tanager ^{atl, Fo}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
Fringillidae (6)			
<i>Spinus magellanicus</i> (Vieillot, 1805)	Hooded Siskin *	ALSP, CSP, Patch	1, 3, 4
<i>Euphonia chlorotica</i> (Linnaeus, 1766)	Purple-throated Euphonia ^{Fo}	ALSP, CSP, Patch	1, 2, 3, 4, 6
<i>Euphonia violacea</i> (Linnaeus, 1758)	Violaceous Euphonia ^{Fo}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
<i>Euphonia cyanocephala</i> (Vieillot, 1818)	Golden-rumped Euphonia ^{Fo}	CSP	1, 6
<i>Euphonia pectoralis</i> (Latham, 1801)	Chestnut-bellied Euphonia ^{atl, Fo}	ALSP, CSP, Patch	1, 2, 3, 4, 5, 6
<i>Chlorophonia cyanea</i> (Thunberg, 1822)	Blue-naped Chlorophonia ^{Fo}	CSP, Patch	1, 2, 3, 4, 6
Estrildidae (1)			
<i>Estrilda astrild</i> (Linnaeus, 1758)	Common Waxbill ^{Exo}	ALSP, CSP, Patch	4, 6
Passeridae (1)			
<i>Passer domesticus</i> (Linnaeus, 1758)	House Sparrow ^{Exo *}	CSP, Patch	1, 4

APPENDIX 2

Species reported for Serra da Cantareira and pending confirmation.

* = Birds pending confirmation according to Graham (1992); ‡ = Species only reported in Figueiredo and Loo (2000) and without any further mention, possibly escaped or intentionally released from captivity; CSP = Cantareira State Park; ALSP = Alberto Löfgren State Park.

Species	English Name	Locality
<i>Crypturellus parvirostris</i> *	Small-billed Tinamou	CSP
<i>Elanus leucurus</i> *	White-tailed Kite	ALSP
<i>Aratinga auricapillus</i> ‡	Golden-capped Parakeet	CSP
<i>Amazona vinacea</i> ‡	Vinaceous Parrot	Urban area near CSP
<i>Hymenops perspicillatus</i> *	Spectacled Tyrant	ALSP
<i>Cyanocorax caeruleus</i> *	Azure Jay	CSP
<i>Gnorimopsar chopi</i> *	Chopi Blackbird	CSP
<i>Sporophila angolensis</i> *	Chestnut-bellied Seed-Finch	CSP
<i>Cyanoloxia brissonii</i> *	Ultramarine Grosbeak	CSP

Submitted: 28 June 2016

Received in revised form: 12 January 2017

Accepted: 21 January 2017

Editorial responsibility: Walter A.P. Boeger

Author Contributions: VRT, MAR and LFS designed sample design, VRT, MAR, ACL, PFD, FS and LFS collected the data, VRT analyzed the data, elaborated figures and wrote the paper, VRT, MAR and LFS revised the manuscript.

Competing Interests: The authors have declared that no competing interests exist.